

DISCOVERY OF X-RAY EMISSION FROM SUPERNOVA 1970G WITH *Chandra*: FILLING THE VOID BETWEEN SUPERNOVAE AND SUPERNOVA REMNANTS

STEFAN IMMLER¹ AND K. D. KUNTZ²

Exploration of the Universe Division, X-Ray Astrophysics Laboratory, Code 662,
 NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

Draft version February 2, 2008

ABSTRACT

We report on the discovery of X-ray emission from SN 1970G in M101, 35 years after its outburst, using deep X-ray imaging with the *Chandra* X-ray observatory. The *Chandra* ACIS spectrum shows that the emission is soft ($\lesssim 2$ keV) and characteristic for the reverse shock region. The X-ray luminosity, $L_{0.3-2} = (1.1 \pm 0.2) \times 10^{37}$ ergs s⁻¹, is likely caused by the interaction of the supernova (SN) shock with dense circumstellar matter. If the material was deposited by the stellar wind from the progenitor, a mass-loss rate of $\dot{M} = (2.6 \pm 0.4) \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ ($v_w/10 \text{ km s}^{-1}$) is inferred. Utilizing the high-resolution *Chandra* ACIS data of SN 1970G and its environment, we reconstruct the X-ray lightcurve from previous *ROSAT* HRI, PSPC, and *XMM-Newton* EPIC observations, and find a best-fit linear rate of decline of $L \propto t^{-s}$ with index $s = 1.7 \pm 0.6$ over a period of 12–35 years after the outburst. As the oldest SN detected in X-rays, SN 1970G allows, for the first time, direct observation of the transition from a SN to its supernova remnant (SNR) phase.

Subject headings: stars: supernovae: individual (SN 1970G) — stars: circumstellar matter — galaxies: individual (M101, NGC 5457) — ISM: supernova remnants — X-rays: general — X-rays: individual (SN 1970G, M101, NGC 4457) — X-rays: ISM

1. INTRODUCTION

Since the launch of the *Chandra* and *XMM-Newton* X-ray observatories, the number of supernovae (SNe) detected in their near aftermath has more than doubled (see Immler & Lewin 2003 for a review article)³. The high-quality X-ray spectra have confirmed the validity of the circumstellar interaction models (see Fransson, Lundqvist & Chevalier 1996 and references therein) which predict a hard spectral component ($\gtrsim 10$ keV) for the forward shock emission during the early epoch ($\lesssim 100$ days) and a soft thermal component ($\lesssim 1$ keV) for the reverse shock emission after the expanding shell has become optically thin. The soft emission component dominates the X-ray output of the interaction regions due to its higher emission measure and higher electron densities. This expected ‘softening’ of the X-ray spectrum has been observationally confirmed for a number of young SNe, such as SNe 1978K (Schlegel et al. 2004), 1979C (Immler et al. 2005), 1993J (Zimmermann & Aschenbach 2003), 1999em and 1998S (Pooley et al. 2002).

Where sufficient data are available, X-ray lightcurves could be established over significant time scales (e.g., ≈ 25 years for SNe 1978K and 1979C; Schlegel et al. 2004, Immler et al. 2005; ≈ 8 years for 1993J; Immler et al. 2001, Zimmermann & Aschenbach 2003). However, little is known about the transition from a young SN into its supernova remnant (SNR) phase as there have been only few detections of intermediate-age (25–300 yrs) SNe in the radio and none in X-rays. While the X-ray emission

of SNe is dominated by the interaction of the shock with the ambient circumstellar matter (CSM), likely deposited by the progenitor’s stellar wind, the emission from a SNR is thought to originate in the shocked interstellar medium (ISM). At an age of ≈ 35 years, SN 1970G is the oldest SN detected in X-rays and closes this gap which allows one, for the first time, to witness the transition from a SN to its SNR phase.

In this paper we report on the *Chandra* X-ray observation of SN 1970G, as well as previous X-ray data from *ROSAT* and *XMM-Newton*. In §2 we briefly describe the data and analysis thereof and report on the X-ray spectrum and multi-mission X-ray lightcurve in §3. We discuss the results in the context of the CSM interaction model and put SN 1970G in the context of the evolution from a SN to a SNR in §4, followed by a summary in §5.

2. DATA PROCESSING AND ANALYSIS

SN 1970G has been observed with the *Chandra* observatory as part of the 1 Ms observation of the host galaxy M101 (NGC 5457; Kuntz 2005). Five individual *Chandra* ACIS observations (sequence numbers 600389, 600389, 600389, 600389) from July 5–11, 2004, were merged into a single observation resulting in a cleaned and exposure-corrected on-source exposure time of 139.5 ks for the ACIS-S2 chip. Details of the data calibration and analysis are explained in Kuntz (2005). The high-resolution ($\approx 0''.5$ FWHM) *Chandra* data were used to search for X-ray emission from the position of SN 1970G, and to examine the contamination by nearby X-ray sources. A *Chandra* ACIS-S2 image of SN 1970G and its environment is given in Fig. 1.

We further constructed the long-term X-ray lightcurve of SN 1970G by extracting source counts from previous *ROSAT* HRI (May 14 – Nov 23, 1996; sequence numbers RH600820N00, RH600820A01; merged expo-

¹ Universities Space Research Association, Code 662, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

² The Henry A. Rowland Department of Physics and Astronomy, The Johns Hopkins University, 3400 Charles Street, Baltimore, MD 21218, USA

³ A complete list of X-ray SNe and references are available at http://lheawww.gsfc.nasa.gov/users/immler/supernovae_list.html




FIG. 1.— Full (0.3–8 keV) band *Chandra* ACIS-S2 image of the region (size $1' \times 1'$) around the position of SN 1970G. The image is smoothed with a Gaussian of $2''.5$ (FWHM) and is given in logarithmic gray-scale ranging from 0 – 1.5 cts pix^{-1} (1 pix corresponds to $0''.5$). Contour lines are 0.2, 0.4 (black), 0.6, 0.8, 1, 1.2, 1.4, and 1.6 cts pix^{-1} (white). The radio position of SN 1970G (Cowan, Goss & Sramek 1991) north-east of the HII region NGC 5455 is marked by a white cross.

sure time 176.1 ks) and *ROSAT* PSPC observations (Jun 8–9, 1991; sequence number RP600108N00; exposure 34.5 ks). Archival *XMM-Newton* EPIC observations from June 4, 2002 (obs-id 0104260101, 43.3 ks) July 23, 2004 (obs-id 0164560701, 41.8 ks) January 8, 2005 (obs-id 0212480201, 32.4 ks) were also analyzed according to standard analysis procedures described in the *XMM-Newton* ABC Guide version 2.01⁴.

A short (2.8 ks) *Einstein* HRI observation 154 days after the outburst of SN 1970G did not result in the detection of the SN and is not included in this study. Archival *ASCA* observations were not used due to the large aperture ($\approx 1'$ FWHM) and contamination by nearby X-ray sources. The *ROSAT* and *XMM-Newton* count rates were further compared to the published results by Wang, Immmler & Pietsch (1999; *ROSAT*) and Jenkins et al. (2005; *XMM-Newton*).

3. RESULTS

A point-like X-ray source is detected at the radio position of SN 1970G (Cowan, Goss & Sramek 1991), with an offset ($\lesssim 0''.5$) similar to the spatial resolution and astrometry of the *Chandra* observation (see Fig. 1). SN 1970G is spatially resolved and well separated from a nearby ($\approx 5''$) HII region (NGC 5455). Exposure, background, and aperture corrected counts extracted from the position of SN 1970G gives a count rate of $(4.3 \pm 0.6) \times 10^{-4}$ cts s^{-1} . While the limited photon statistics does not allow a detailed characterization of the energy distribution of the recorded photons, spectral analysis shows that the emission is soft, with the bulk of

the emission being confined to the $\lesssim 2$ keV band.

Adopting a $kT = 0.6$ keV thermal plasma spectrum typical for the late-time ($\gtrsim 100$ days) emission of SNe and assuming a Galactic foreground column density with no intrinsic absorption ($N_{\text{H}} = 1.16 \times 10^{20} \text{ cm}^{-2}$; Dickey & Lockman 1990) gives a 0.3–2 keV flux and luminosity of $f_{0.3-2} = (1.8 \pm 0.3) \times 10^{-15} \text{ ergs cm}^{-2} \text{ s}^{-1}$ and $L_{0.3-2} = (1.1 \pm 0.2) \times 10^{37} \text{ ergs s}^{-1}$, respectively, for a distance of 7.2 Mpc (Stetson et al. 1998).

Since no information is available about the spectral evolution of SN 1970G or other SNe at this age, we used the same spectral template to convert *ROSAT* and *XMM-Newton* count rates into fluxes and luminosities to construct the long-term (12–35 years) X-ray lightcurve.

An X-ray source is visible in the *ROSAT* HRI images of M101 at the position of SN 1970G (see Fig. 2 in Wang, Immmler & Pietsch 1999). The source, however, is not included in the *ROSAT* HRI catalog since it is slightly below the employed detection threshold ($S/N = 3.5$; Wang, Immmler & Pietsch 1999). Extraction of source counts within the HRI aperture gives a 0.1–2.4 keV count rate of $(2.9 \pm 0.6) \text{ cts s}^{-1}$, corresponding to a 0.3–2 keV band flux and luminosity of $f_{0.3-2} = (6.7 \pm 1.4) \times 10^{-15} \text{ ergs cm}^{-2} \text{ s}^{-1}$ and $L_{0.3-2} = (4.2 \pm 0.9) \times 10^{37} \text{ ergs s}^{-1}$, respectively.

An X-ray source is also listed in the *ROSAT* PSPC catalog at the position of SN 1970G (source P12; Wang, Immmler & Pietsch 1999), with a signal-to-noise ratio of $S/N = 4.5$. Given that no other X-ray source is visible in the high-resolution *Chandra* image in the environment of SN 1970G within the PSPC aperture ($\approx 25''$ FWHM), except for low-level emission of a neighboring HII region (NGC 5455, see Fig. 1), it is safe to assume that the PSPC source is due to the SN itself. Assuming the emission arises from SN 1970G, a (0.5–2 keV band) count rate of $(1.3 \pm 0.1) \times 10^{-3} \text{ cts s}^{-1}$ is inferred, corresponding to $f_{0.3-2} = (1.3 \pm 0.1) \times 10^{-14} \text{ ergs cm}^{-2} \text{ s}^{-1}$ and $L_{0.3-2} = (8.4 \pm 0.9) \times 10^{37} \text{ ergs s}^{-1}$.

A faint X-ray source is visible in each of the nine *XMM-Newton* EPIC images at low energies (< 2 keV). No photons were recorded in the hard (2–6 keV) band. The source, however, is extended ($\approx 20''$) and shows no concentration of the recorded photons at the radio position of SN 1970G. Since it is likely that the emission is contaminated by the nearby HII region and other soft emission components inside the galactic disk, we only use the count rates, fluxes, and luminosities as upper limits. Discrepancies between the published PN and MOS count rates for source #44 (factor $\times 16.5$; Jenkins et al. 2005) for one of the published *XMM-Newton* data (obs-id 0104260101) makes a comparison problematic.

We calculated the mass-loss rate of the progenitor as a function of the stellar wind age using the relationship $L_{\text{x}} = 4/(\pi m^2) \Lambda(T) \times (\dot{M}/v_{\text{w}})^2 \times (v_{\text{s}} t)^{-1}$, where m is the mean mass per particle ($2.1 \times 10^{-27} \text{ kg}$ for a H+He plasma), $\Lambda(T)$ the cooling function of the heated plasma at temperature T , $\dot{M}$ the mass-loss rate of the progenitor, v_{w} the speed of the stellar wind blown off by the progenitor, and v_{s} the speed of the outgoing shock (see Immmler, Wilson & Terashima 2002). A constant shock velocity of $v_{\text{s}} = 9,000 \text{ km s}^{-1}$ similar to other core-collapse SNe, such as SN 1979C (Marcaide et al. 2002) was assumed. An effective (0.3–2 keV band) cooling function

⁴ available from <http://heasarc.gsfc.nasa.gov/docs/xmm/abc/>

TABLE 1
X-RAY PROPERTIES OF SN 1970G

Epoch [years]	Instrument	Count Rate [10^{-4} cts s^{-1}]	$f_{0.3-2}$ [10^{-15} ergs cm^{-2} s^{-1}]	$L_{0.3-2}$ [10^{37} ergs s^{-1}]	$\dot{M}$ [10^{-4} $M_{\odot}$ yr^{-1}]
11.9	<i>ROSAT</i> PSPC	13.0 ± 1.3	13.4 ± 1.4	8.4 ± 0.9	4.2 ± 0.4
16.8	<i>ROSAT</i> HRI	2.9 ± 0.6	6.7 ± 1.4	4.2 ± 0.9	3.4 ± 0.7
31.9	<i>XMM-Newton</i> EPIC	< 48.3	< 10.2	< 6.3	< 5.8
34.0	<i>XMM-Newton</i> EPIC	< 80.6	< 17.0	< 10.6	< 7.8
34.5	<i>XMM-Newton</i> EPIC	< 119.3	< 25.2	< 15.6	< 9.4
34.7	<i>Chandra</i> ACIS	4.3 ± 0.6	1.8 ± 0.3	1.1 ± 0.2	2.6 ± 0.4

of $\Lambda = 3 \times 10^{-23}$ ergs cm^3 s^{-1} for an optically thin thermal plasma with a temperature of 10^7 K was adopted, which corresponds to the assumed thermal plasma temperature. Assuming different plasma temperatures in the range 0.5–1 keV would lead to changes in the emission measure of $\lesssim 10\%$. Key observational properties of SN 1970G are listed in Table 1.

4. DISCUSSION

Previous X-ray observations of SN 1970G with *ROSAT*, *ASCA*, and *XMM-Newton* lacked the spatial resolution needed to separate the SN from the nearby ($\approx 5''$) HII region NGC 5455. Using the sub-arcsec *Chandra* imaging capability, in combination with the deep exposure time (140 ks), we detect soft ($\lesssim 2$ keV) X-ray emission from SN 1970G with a luminosity of $L_{0.3-2} \approx 1 \times 10^{37}$ ergs s^{-1} , 35 years after its outburst. The source is spatially resolved from the nearby HII region NGC 5455 ($\approx 5''$), which is observed at a $\lesssim 20\%$ flux level compared to SN 1970G. Contamination of the previous *ROSAT* data of SN 1970G with the HII region should therefore be minimal, especially when kept in mind that at earlier epochs SN 1970G was significantly brighter, while the HII region presumably has a constant flux.

Given the information from the high-resolution spatial imaging with *Chandra*, we recover X-rays from SN 1970G in the *ROSAT* HRI and PSPC observations. SN 1970G is not conclusively detected in three recent (2002–2005) *XMM-Newton* observations due to the extent of the source ($\approx 20''$) as imaged by the PN and MOS instruments, and positional offsets larger than the instrumental uncertainties ($\approx 5''$).

In combination, an X-ray rate of decline of $L \propto t^{-s}$ with index $s = 1.7 \pm 0.6$ is similar to other X-ray SNe (e.g., SN 1999em, Pooley et al. 2002; SN 1986J, Houck et al. 1998). This strengthens the case that the emission is due to the SN itself, even though a variable source in the environment of NGC 5455 cannot be entirely ruled out. The lack of X-rays above 2 keV confirms that the emission is soft and of possible thermal original. Such a soft emission is expected for the late emission of a SN, where the shocked emission originates from the reverse shock region.

Assuming that the emission arises from shock heated plasma deposited by the progenitor's stellar wind, a mass-loss rate of $\dot{M} = (2.6 \pm 0.4) \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ ($v_w/10 \text{ km s}^{-1}$) is inferred for the late epoch observed with *Chandra*. A similar mass-loss rate was inferred from radio observations [$2 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ ($v_w/10 \text{ km s}^{-1}$); Weiler 1993]. The

mass-loss rates for the earlier epoch are slightly higher and might indicate an evolution in the mass-loss rate history of the progenitor (see Tab. 1). It should be noted, however, that the lack of knowledge about the shock front velocity precludes a more detailed study. If the shock front experienced a significant deceleration, as expected when the shock has swept up a significant amount of CSM, the lower inferred mass-loss rates for the later epochs would be even lower (see the discussion about the mass-loss rate history of SN 1979C in Immler et al. 2005). However, given the uncertainties in the derived X-ray luminosities due to a possible residual contamination of the *ROSAT* and *XMM-Newton* data caused by their larger point-spread functions (PSF; $\approx 5''$ FWHM for the *ROSAT* HRI; $5''$ – $6''$ FWHM for the *XMM-Newton* EPIC instrument; $\approx 25''$ FWHM for the *ROSAT* PSPC), the apparent increase of the mass-loss rate at earlier epochs is not regarded as significant. Interestingly, the mass-loss rates are higher for the instruments with the largest PSF (EPIC and PSPC). If the mass-loss rates inferred from the instruments with the smallest PSF are compared [$\dot{M} = (3.4 \pm 0.7)$ and $(2.6 \pm 0.4) \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ ($v_w/10 \text{ km s}^{-1}$) for the HRI and ACIS, respectively], no statistically significant differences are observed. We therefore conclude that SN 1970G shows a mass-loss rate which is consistent with being constant over a period of 12–35 years after the peak optical brightness (July 30, 1970; Detre 1970), corresponding to $\approx 11,000$ – $31,000$ years in the stellar wind age using the conversion $t_w = tv_s/v_w$ and assuming a stellar wind speed of 10 km s^{-1} and a shock front velocity of $9,000 \text{ km s}^{-1}$.

The measured mass-loss rate for SN 1970G is similar to those inferred for other Type II SNe, which typically range from 10^{-5} to $10^{-4} M_{\odot} \text{ yr}^{-1}$ (see Immler & Lewin 2003). This is indicative that the X-ray emission arises from shock-heated CSM deposited by the progenitor rather than shock-heated ISM, even at this late epoch after the outburst.

In order to explain the X-ray luminosities of SNRs such as Cas A ($L_{0.3-2} = 2.8 \times 10^{37}$ ergs s^{-1} at an age of ≈ 320 years), mass-loss rates in excess of $10^{-4} M_{\odot} \text{ yr}^{-1}$ would be required in the shocked stellar wind scenario. The high required amounts of deposited material could not be provided by a sustained stellar wind since the wind would interact with the ambient ISM at these large radii ($\gtrsim 10^{18} \text{ cm}$) from the site of the explosion. Instead, the X-ray emission from SNRs is thought to originate in radiative cooling of the shock-heated ISM which has substantially larger densities. The relatively low inferred




FIG. 2.— Soft (0.3–2 keV) band X-ray luminosities of all SNe detected to date (filled squares) and historical SNRs (filled triangles) as a function of age (in units of years). The X-ray lightcurve of SN 1970G is marked by filled diamonds with error bars (from left to right: *ROSAT* PSPC, *ROSAT* HRI and *Chandra* ACIS). *XMM-Newton* EPIC upper limits are indicated by arrows.

mass-loss rate required to produce the observed X-ray luminosity of SN 1970G indicates that the SN shock has not yet reached the ISM. Consequently, the SN 1970G has not yet reached its SNR phase.

We therefore conclude that at an age of 35 years, SN 1970G is the oldest SN observed in X-rays, and provides important constraint as to when the transition from a SN to its SNR phase occurs, which is likely at a later stage (≈ 50 – 100 years) after its outburst.

In Fig. 2 we plot the X-ray evolution of all SNe detected to date⁵ as well as the X-ray luminosities of historical Galactic, LMC, SMC (as observed with *Chandra*⁶), M31 (Supper et al. 2001; Kong et al. 2003), and M33 (Ghavamian et al. 2005) SNRs. A comparison of the X-ray luminosities shows that the transition from a SN to a SNR appears to be rather smooth. If the evolution of SN 1970G continues as observed to date, it will be similar to X-ray faint Galactic SNRs such as SNRs 1181 and 1006. X-ray bright SNe such as SNe 1978K and 1979C, which show no sign for an X-ray evolution out to large distances ($\approx 10^{17}$ cm) at an age of ≈ 25 years after their outburst, can evolve into luminous SNRs such as Cas A

due to their strong interaction with their dense ambient CSM, or possibly attain the X-ray luminosities of the brightest SNRs observed (such as W49B or N132D) in case of a dense ISM interaction.

Clearly, X-ray detections and monitoring of intermediate-age (around 100 years) SNe are needed to address these questions and to study the transition from a SN into a SNR in detail. Deep galaxy surveys with *Chandra* and *XMM-Newton* and future X-ray missions such as *Const-X* and *XEUS* provide the capabilities to detect these intermediate-age SNe in the near future and to fill the void in the evolution of a SN to a SNR.

5. SUMMARY

Based on our deep (140 ks) X-ray imaging of M101 with the *Chandra* observatory we present the detection of SN 1970G in X-rays. Key results are:

- SN 1970G is detected with a luminosity of $L_{0.3-2} = (1.1 \pm 0.2) \times 10^{38}$ ergs s⁻¹, 35 years after its outburst in our deep *Chandra* observation of M101.
- SN 1970G is recovered in previous *ROSAT* HRI and PSPC observations and shows a best-fit linear rate of decline of $L \propto t^{-s}$ with index $s = 1.7 \pm 0.6$ over the observed period of 12–35 yrs after its outburst, consistent with recent *XMM-Newton* observations.
- The emission of SN 1970G is soft ($\lesssim 2$ keV) and indicates that it originates in the reverse shocked region, as expected for the late emission in the evolution of a SN.
- A mass-loss rate of $\dot{M} = (2.6 \pm 0.4) \times 10^{-5} M_{\odot} \text{ yr}^{-1}$ is inferred consistent with being constant over a period of 11,000–31,000 years in the evolution of the progenitor.
- At an age of 35 years, the SN has not yet reached its SNR phase, which is likely to occur at a later stage in the evolution (≈ 50 – 100 years).
- Comparison of the X-ray lightcurves of all SNe detected to date with the X-ray luminosities of SNRs indicates a smooth transition of SNe into their SNR phases.

This paper is based on observations obtained with *XMM-Newton*, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA. KDK was supported by the *Chandra* grant SAO GO-5600587. The authors wish to thank R. Petre for helpful suggestions and comments.

⁵ http://lheawww.gsfc.nasa.gov/users/immler/supernovae_list.html

⁶ <http://www.astro.psu.edu/users/green/Main/main5.html>

REFERENCES

- Cowan, J. J., Goss, W. M., Sramek, R. A. 1991, *ApJ*, 379, L49
 Detre, L. 1970, *IAU Circ.*, No. 2269
 Dickey, J. M., Lockman, F. J. 1990, *ARAA*, 28, 215
 Fransson, C., Lundqvist, P., Chevalier, R. A. 1996, *ApJ*, 461, 993
 Ghavamian, P., Blair, W. P., Long, K. S. et al. 2005, *AJ*, submitted (astro-ph/0504583)
 Houck, J. C., Bregman, J. N., Chevalier, R. A., Tomisaka, K. 1998, *ApJ*, 493, 431
 Immler, S., Aschenbach, B., Wang, Q. D. 2001, *ApJ*, 561, L107
 Immler, S., Wilson, A. S., Terashima, Y. 2003, *ApJ*, 573, L27
 Immler, S., Lewin, W. 2003, in *Supernovae and Gamma-Ray Bursters* (ed K. Weiler), *Lecture Notes in Physics*, 598, 91
 Immler, S., Fesen, R. A., Van Dyk, S. D. et al. 2005, *ApJ*, submitted (astro-ph/0503678)
 Jenkins, L. P., Roberts, T. P., Warwick, R. S., Kilgard, R. E., Ward, M. J. 2005, *MNRAS*, 357, 401
 Kong, A. K. H., DiStefano, R., Garcia, M. R., Greiner, J. 2003, *ApJ*, 585, 298
 Kuntz, K. 2005, *ApJ*, in prep.
 Marcaide, J. M., Pérez-Torres, M. A., Ros, E., Alberdi, A., Diamond, P. J., Guirado, J. C., Lara, L., Van Dyk, S. D., Weiler, K. W. 2002, *A&A*, 384, 408
 Pooley, D., Lewin, W. H. G., Fox, D. W., et al. 2002, *ApJ*, 572, 932
 Schlegel, E. M., Kong, A., Kaaret, P., DiStefano, R., Murray, S. 2004, *ApJ*, 603, 644
 Stetson, P. B., Saha, A., Ferrarese, L. et al. 1998, *ApJ*, 508, 491
 Supper, R., Hasinger, G., Lewin, W. H. G. 2001, *A&A*, 373, 63

- Weiler, K. W. 1993, in ASP Conference Series 'Massive Stars: Their Lives in the Interstellar Medium' (eds J.P. Cassinelli, E.B. Churchwell), Vol. 35, 436
- Wang, Q. D., Immler, S., Pietsch, W. 1999, ApJ, 523, 121
- Zimmermann, H.-U., Aschenbach, B. 2003, A&A, 406, 969